



Norges teknisk-naturvitenskapelige universitet
Institutt for matematiske fag

TMA4285
Tidsrekkemodeller

Løsningsforslag - Øving 3, 2005

Exercise 2.8

$$X_t = \phi X_{t-1} + Z_t$$

$$\begin{aligned} X_t &= \phi X_{t-1} + Z_t \\ &= Z_t + \phi(Z_{t-1} + \phi X_{t-2}) \\ &= \dots \\ &= Z_t + \phi Z_{t-1} + \dots + \phi^n Z_{t-n} + \phi^{n+1} X_{t-n-1} \end{aligned}$$

That is

$$X_t - \phi^{n+1} X_{t-n-1} = Z_t + \phi Z_{t-1} + \dots + \phi^n Z_{t-n}$$

First we calculate

$$\begin{aligned} \text{Var}(X_t - \phi^{n+1} X_{t-n-1}) &= \gamma(0)(1 + \phi^{2n+2}) - 2\phi^{n+1}\gamma(n+1) \\ &\leq \gamma(0)(1 + |\phi|^{2n+2} + 2|\phi|^{n+1}) = 4\gamma(0) \end{aligned}$$

if X_t is stationary and $|\phi| = 1$

Next we calculate

$$\text{Var}(Z_t + \phi Z_{t-1} + \dots + \phi^n Z_{t-n}) = n\sigma^2$$

if $|\phi| = 1$

Since clearly $n\sigma^2 > 4\gamma(0)$ for sufficiently large n , we have reached a contradiction. Hence X_t cannot be stationary if $|\phi| = 1$.

Exercise 2.10

$$X_t - \phi X_{t-1} = Z_t + \theta Z_{t-1}$$

where $\phi = \theta = 0.5$

According to Section 2.3, equation (2.3.3), we obtain that

$$X_t = \sum_{j=0}^{\infty} \psi_j Z_{t-j}$$

where $\psi_0 = 1$, $\psi_j = (\phi + \theta)\phi^{j-1} = 0.5^{j-1}$ for $j = 1, 2, \dots$

From Section 2.3, equation (2.3.5), we get

$$Z_t = \sum_{j=0}^{\infty} \pi_j X_{t-j}$$

where $\pi_0 = 1$, $\pi_j = -(\phi + \theta)(-\theta)^{j-1} = -(-0.5)^{j-1}$ for $j = 1, 2, \dots$

Agrees with the results from ITSM.

Exercise 2.12

The given MA(1)-model is

$$X_t = Z_t - 0.6Z_{t-1}$$

where $Z_t \sim \text{WN}(0, 1)$.

Observed that $\bar{x}_{100} = 0.157$

The variance of \bar{x}_{100} :

$$\begin{aligned} \text{Var}[\bar{x}_{100}] &= \frac{1}{n} \sum_{h=-n}^n \left(1 - \frac{|h|}{n}\right) \gamma(h) \\ &= \frac{1}{100} \left(\gamma(0) + 2 \cdot \frac{99}{100} \gamma(1) \right) \\ &= \frac{1}{100} (1.36 - 1.98 \cdot 0.6) \\ &= 0.00172 \end{aligned}$$

That is, 95% confidence bounds for μ are approximately

$$\begin{aligned} &\bar{x}_{100} \pm 1.96\sqrt{0.00172} \\ &= 0.157 \pm 1.96 \cdot 0.0415 = 0.157 \pm 0.0813 = 0.076, 0.238 \end{aligned}$$

Reject $H_0: \mu = 0$ in favour of the alternative hypothesis $H_1: \mu \neq 0$ at significance level 0.05 since the 95% bounds for μ do not include the value 0.

Note: The conclusion would differ if the time series $X_t \sim \text{IID}(0, 1.36)$.

Exercise 2.13

a)

Assume an AR(1)-model

$$X_t = \phi X_{t-1} + Z_t$$

Since $\rho(h) = \phi^h$ ($h > 0$) for an AR(1)-model, and it has been observed that $\hat{\rho}(2) = 0.145$, we shall assume that $\phi^2 \ll 1$. Using Bartlett's formula, the following approximate relations are obtained:

$$\text{Var}[\hat{\rho}(1)] \approx \frac{1}{n}(1 - \phi^2)$$

and

$$\text{Var}[\hat{\rho}(2)] \approx \frac{1}{n}(1 - \phi^2)(1 + 3\phi^2)$$

That is, 95% confidence bounds for $\rho(1)$ are approximately

$$\hat{\rho}(1) \pm \frac{1.96}{\sqrt{n}} \sqrt{1 - \phi^2}$$

Correspondingly, 95% confidence bounds for $\rho(2)$ are approximately

$$\hat{\rho}(2) \pm \frac{1.96}{\sqrt{n}} \sqrt{(1 - \phi^2)(1 + 3\phi^2)}$$

With $\phi = \hat{\phi} = \hat{\rho}(1)$, $n = 100$, $\hat{\rho}(1) = 0.438$, $\hat{\rho}(2) = 0.145$, these bounds become for $\rho(1)$: 0.262, 0.614, and for $\rho(2)$: -0.073, 0.369.

These values are not consistent with $\phi = 0.8$, since both $\rho(1) = 0.8$ and $\rho(2) = 0.64$ are outside these bounds.

b)

Assume an MA(1)-model

$$X_t = Z_t + \theta Z_{t-1}$$

Bartlett's formula gives the following approximate relations

$$\text{Var}[\hat{\rho}(1)] \approx \frac{1}{n}(1 - 3\rho(1)^2 + 4\rho(1)^4)$$

and

$$\text{Var}[\hat{\rho}(2)] \approx \frac{1}{n}(1 + 2\rho(1)^2)$$

That is, 95% confidence bounds for $\rho(1)$ are approximately

$$\hat{\rho}(1) \pm \frac{1.96}{\sqrt{n}} \sqrt{1 - 3\rho(1)^2 + 4\rho(1)^4}$$

Correspondingly, 95% confidence bounds for $\rho(2)$ are approximately

$$\hat{\rho}(2) \pm \frac{1.96}{\sqrt{n}} \sqrt{1 + 2\rho(1)^2}$$

With the numbers as in a), it is now obtained that these bounds become for $\rho(1)$: 0.290, 0.586, and for $\rho(2)$: -0.082, 0.378.

$\theta = 0.6$ leads to $\rho(1) = \frac{\theta}{1+\theta^2} = 0.4412$, $\rho(2) = 0$. It follows that the confidence bounds are consistent with these two values, and the data are therefore consistent with the MA(1)-model $X_t = Z_t + 0.6Z_{t-1}$

Exercise 2.14

$$X_t = A \cos(\omega t) + B \sin(\omega t), \quad t \in \mathbb{Z}$$

where A and B are uncorrelated random variables with zero mean and variance 1. This process is stationary with ACF $\rho(h) = \cos(\omega h)$.

a)

$$P_1 X_2 = \phi_{11} X_1$$

where $\gamma(0)\phi_{11} = \gamma(1)$, which gives $\phi_{11} = \rho(1) = \cos \omega$. Hence

$$P_1 X_2 = \cos(\omega) X_1$$

Also

$$E[(X_2 - P_1 X_2)^2] = \gamma(0) - \phi_{11}\gamma(1) = \gamma(0)(1 - \cos^2 \omega) = \sin^2 \omega$$

Note: 2.14 is an example in which the matrix Γ_n in the equation $\Gamma_n \bar{\phi}_n = \bar{\gamma}_n$ is singular for $n \geq 3$. This is because $X_3 = (2 \cos \omega) X_2 - X_1$.

b)

$$P_2 X_3 = \phi_{21} X_2 + \phi_{22} X_1$$

where

$$\begin{aligned}\gamma(0)\phi_{21} + \gamma(1)\phi_{22} &= \gamma(1) \\ \gamma(1)\phi_{21} + \gamma(0)\phi_{22} &= \gamma(2)\end{aligned}$$

that is

$$\begin{aligned}\phi_{21} + (\cos \omega)\phi_{22} &= \cos \omega \\ (\cos \omega)\phi_{21} + \phi_{22} &= \cos 2\omega\end{aligned}$$

Solving these equations give $\phi_{22}(\cos^2 \omega - 1) = \cos^2 \omega - 2 \cos^2 \omega + 1 = -\cos^2 \omega + 1$, that is, $\phi_{22} = -1$, and then, $\phi_{21} = \cos \omega - \phi_{22} \cos \omega = 2 \cos \omega$. Hence

$$P_2 X_3 = (2 \cos \omega) X_2 - X_1$$

and

$$\begin{aligned}E[(X_3 - P_2 X_3)^2] &= \gamma(0) - \bar{\phi}_2 \bar{\gamma}_2 \\ &= 1 - (2 \cos \omega, -1)(\cos \omega, \cos 2\omega) \\ &= 1 - 2 \cos^2 \omega + \cos 2\omega = 0\end{aligned}$$

c)

From b) and stationarity, it follows that

$$P(X_{n+1}|X_n, X_{n-1}) = (2 \cos \omega)X_n - X_{n-1}$$

with $\text{MSE} = 0$.

Since $(2 \cos \omega)X_n - X_{n-1}$ is a linear combination of X_s , $-\infty < s \leq n$, and since it is impossible to find a predictor of this form with smaller MSE, we conclude that $\tilde{P}_n X_{n+1} = (2 \cos \omega)X_n - X_{n-1}$ with $\text{MSE} = 0$.